



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

10/511096

Rec'd PCT/PTO 14 OCT 2004

REC'D 27 MAY 2003

WIPO

PCT

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-
gen stimmen mit der
ursprünglich eingereichten
Fassung der auf dem näch-
sten Blatt bezeichneten
europäischen Patentanmel-
dung überein.

The attached documents
are exact copies of the
European patent application
described on the following
page, as originally filed.

Les documents fixés à
cette attestation sont
conformes à la version
initialement déposée de
la demande de brevet
européen spécifiée à la
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

02008553.6

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk



Anmeldung Nr:
Application no.: 02008553.6
Demande no:

Anmeldetag:
Date of filing: 16.04.02
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

EVOTEC Neurosciences GmbH
Schnackenburgallee 114
22525 Hamburg
ALLEMAGNE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Diagnostic and therapeutic use of a Golgi protein for neurodegenerative diseases

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

G01N33/48

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE TR

DIAGNOSTIC AND THERAPEUTIC USE OF A GOLGI PROTEIN FOR NEURODEGENERATIVE DISEASES

The present invention relates to methods of diagnosing, prognosticating and monitoring the progression of neurodegenerative diseases in a subject. Furthermore, methods of therapy control and screening for modulating agents of neurodegenerative diseases are provided. The invention also discloses pharmaceutical compositions, kits, and recombinant animal models.

Neurodegenerative diseases, in particular Alzheimer's disease (AD), have a strongly debilitating impact on a patient's life. Furthermore, these diseases constitute an enormous health, social and economic burden. AD is the most common age-related neurodegenerative condition affecting about 10 % of the population over 65 years of age and up to 45 % over age 85 (for a recent review see Vickers et al., 2000, Progress in Neurobiology, 60:139-165). Presently, this amounts to an estimated 12 million cases in the US, Europe, and Japan. This situation will inevitably worsen with the demographic increase in the number of old people ("aging of the baby boomers") in developed countries. The neuropathological hallmarks that occur in the brains of individuals with AD are senile plaques, composed of amyloid- β protein, and profound cytoskeletal changes coinciding with the appearance of abnormal filamentous structures and the formation of neurofibrillary tangles. AD is a progressive disease that is associated with early deficits in memory formation and ultimately leads to the complete erosion of higher cognitive function. A characteristic feature of the pathogenesis of AD is the selective vulnerability of particular brain regions and subpopulations of nerve cells to the degenerative process. Specifically, the temporal lobe region and the hippocampus are affected early and more severely during the progression of the disease. On the other hand, neurons within the frontal cortex, occipital cortex, and the cerebellum remain largely intact and are protected from neurodegeneration (Terry et al., 1981, Annals of Neurology, 10:184-192).

Currently, there is no cure for AD, nor is there an effective treatment to halt the progression of AD or even to diagnose AD ante-mortem with high probability. Several risk factors have been identified that predispose an individual to develop AD, among them most prominently the epsilon4 allele of apolipoprotein E (ApoE). Although there are rare examples of early-onset AD which have been attributed to genetic defects in the genes for amyloid precursor protein (APP), presenilin-1, and presenilin-2, the prevalent form of late-onset sporadic AD is of hitherto unknown etiologic origin. The late onset and complex pathogenesis of neurodegenerative disorders pose a formidable challenge to the development of therapeutic and diagnostic agents. It is crucial to expand the pool of potential drug targets and diagnostic markers. It is therefore an object of the present invention to provide insight into the pathogenesis of neurodegenerative diseases and to provide methods, materials, and animal models which are suited inter alia for the diagnosis and development of a treatment of these diseases. This object has been solved by the features of the independent claims. The subclaims define preferred embodiments of the present invention.

The Golgi-complex is an intracellular network which was first described in 1898. It has been shown to function as an organelle responsible for the processing, transporting and sorting of intracellular and secreted proteins (reviewed in Nilsson and Warren, 1994, Curr. Opin. Cell Biol, 6:517-521). Localized at the perinuclear site of cells, the Golgi-apparatus can be described as stacks of membranous cisternae which form functionally distinct networks. Briefly, membrane proteins are routed via the endoplasmic reticulum in vesicles through the cis-, medial- and trans-Golgi network and are then transported to their intracellular destination. The transport vesicles which mediate the transport bud from donor membranes and are transported to and fused with an acceptor membrane. The control of these events so far is poorly understood although several proteins have been characterized which play important roles in the targeting and transport of the vesicles, among them being coating proteins (COPs), adaptins, GTP-binding proteins, ADP-ribosylation factors (ARFs), and resident proteins. Several auto-antigens that

are responsible for auto-immune diseases have been shown to be integral parts of the Golgi-apparatus. Such diseases are Sjögren's disease, rheumatoid arthritis or systemic lupus erythematosus (see review by Chan and Fritzler, 1998, *Electr. J. Biotechn.*, 1:1-10). Common to those diseases is the fact that the auto-antigens represent a class of proteins with extended coiled coil domains and non alpha-helical domains at their N- and C-termini. So far, several Golgi auto-antigens are known which are referred to as golgins, such as golgin-95/GM130, golgin-97, golgin-256, golgin-160/GCP170, giantin/macrogolgin/GCP372, and golgin-245/p230. Currently, it is postulated that the golgins form intermolecular complexes that in concert with other proteins serve as docking stations for vesicles and are important for guiding the vesicles through the Golgi-apparatus.

Golgin-245, also referred to as p230, was first identified by antibodies derived from a patient suffering from Sjögren's syndrome (Kooy et al., 1992, *J. Biol. Chem.*, 267:20255-20263). Indirect immunofluorescence analysis revealed that the protein is localized at the Golgi-apparatus, and it has been hypothesized that the protein plays an important role in compartmentalization of the Golgi-apparatus or in sorting and transport of proteins. Subsequently, golgin-245 was cloned and molecularly characterised by two independent groups (Fritzler et al., 1995, *J. Biol. Chem.*, 270:31263-31268; Erlich et al., 1996, *J. Biol. Chem.*, 271:8328-8337). The proteins described in these two studies have been shown to be identical except for an additional 145 amino acids at the N-terminus of the longer isoform. It turned out that the longer isoform of the protein is encoded by an open reading frame of 6690 base pairs and is comprised of 2230 amino acids, resulting in a molecular weight of ~261 kDa (GenBank accession number U41740). Two alternatively spliced mRNAs of approximately 7.7kb have been detected which differ by 21-base pair and 63-base pair inserts in the 3'-region of the gene. The gene coding for golgin-245 has been mapped to chromosome 6p12-22 (Erlich et al., *ibid*). Secondary structure analysis predicts an extraordinary high level of coiled-coil elements, and it has been speculated that these regions might mediate multimerization or the induction of conformational changes as shown for other coiled-coil

proteins. The protein is very hydrophilic and shares a 17-20% homology with other coiled-coil proteins such as kinesin related microtubule motor proteins. In addition, homology has been observed with the granin family of proteins which are present in the secretory granules of neuroendocrine cells (Erich et al., *ibid*).

Golgin-245 has been shown to be associated with vesicles budding from the trans-Golgi network (Gleeson et al., 1996, *J. Cell Sci.*, 109:2811-2821). The protein faces the intracellular compartment and recycles between cytosol and trans-Golgi derived vesicles. Golgin-245 is found primarily on a defined subset of these vesicles and might play a role in the assembly of said vesicles.

The Golgi-targeting sequence has been narrowed down to a stretch of 42 amino acids located at the C-terminus of golgin-245 (Kjer-Nielsen et al., 1999, *J. Cell Sci.*, 112:1645-1654). This domain is highly homologous within the golgin-family of proteins and is characterized by a conserved tyrosine residue within said stretch (Munro and Nichols, 1999, *Curr. Biol.*, 9:377-380). The GRIP-domain has also been shown to bind to rab6, a member of a class of proteins thought to regulate vesicle docking and membrane-tethering (Barr, 1999, *Curr. Biol.*, 9:381-384). The Golgin-family of proteins has only recently been assigned a role in maintaining the structural scaffold which is responsible for the integrity of the Golgi-apparatus (Seeman et al., 2000, *Nature*, 407:1022-1026). According to that study, the golgins can be separated from Golgi-enzymes and are sufficient for a correct rebuilding of the Golgi-apparatus. Hence it is speculated that they may constitute a network by binding either directly or indirectly to the Golgi membranes, implying that the Golgi apparatus functions as an autonomous organelle rather than representing a temporary membranous system being in equilibrium between endoplasmic reticulum and secretory vesicles. Golgin-245 has been found to bind to ADP-ribosylation factor (ARF)-related proteins (ARL) (Van Valkenburgh et al., 2001, *J. Biol. Chem.*, 276:22826-22837). ARL-proteins share a 40-60% identity to ARFs, small GTP-binding proteins. However, ARLs are devoid of enzymatic activities, and it is speculated that they function as binding partners for golgin-245 at the Golgi apparatus.

Golgins are a target for caspases (Mancini et al., 2000, J. Cell Biol., 149:603-612). In a recent report it has been proposed that apoptotic signals may be passed through the Golgi apparatus by the specific cleavage of golgin-160 by caspase-2. Since Golgi autoantigens in patients with systemic auto-immune diseases are frequently cleaved by caspases, and golgin-245 represents the major auto-antigen in Sjögren's disease, it might be speculated that golgin-245 may also play a role in apoptotic signal transduction.

The integrity of intracellular transport processes is a valuable target for the treatment of several disorders, among them neurological and neurodegenerative disorders. It is a feature of the present invention to modulate the interaction of golgin-245 with its target molecules in order to influence processing, trafficking and sorting of intracellular and/or secreted proteins. Of special interest in this context is the fact that one of the key players of Alzheimer's disease, amyloid precursor protein (APP), matures during the secretory pathway through the Golgi apparatus, and it has been speculated that the proteolytic processing of APP, which yields the highly amyloidogenic A β 42, takes place in the trans-Golgi compartment (Greenfield et al., 1999, Proc. Natl. Acad. Sci., 96:742-747). To date, there are no drugs on the market nor in clinical development which specifically and potently target proteins of the golgin family, in particular golgin-245.

In the present invention, using an unbiased and sensitive differential display approach, a transcription product of the gene coding for golgin-245 is detected in human brain samples. Importantly, the present invention discloses an up-regulation of golgin-245 transcripts in the inferior temporal lobe of brain samples taken from AD patients relative to frontal cortex samples. No such up-regulation is observed in samples from age-matched healthy controls. To date, no experiments have been described that demonstrate a relationship between the dysregulation of golgin-245 gene expression and the pathology of neurodegenerative disorders, in particular AD. Such a link, as disclosed in the present invention, offers new ways, inter alia, for the diagnosis and treatment of said disorders, in particular AD.

The singular forms "a", "an", and "the" as used herein and in the claims include plural reference unless the context dictates otherwise. For example, "a cell" means as well a plurality of cells, and so forth. The term "and/or" as used in the present specification and in the claims implies that the phrases before and after this term are to be considered either as alternatives or in combination. For instance, the wording "determination of a level and/or an activity" means that either only a level, or only an activity, or both a level and an activity are determined. The term "level" as used herein is meant to comprise a gage of, or a measure of the amount of, or a concentration of a transcription product, for instance an mRNA, or a translation product, for instance a protein or polypeptide. The term "activity" as used herein shall be understood as a measure for the ability of a transcription product or a translation product to produce a biological effect or a measure for a level of biologically active molecules. The term "activity" also refers to enzymatic activity. The terms "level" and/or "activity" as used herein further refer to gene expression levels or gene activity. Gene expression can be defined as the utilization of the information contained in a gene by transcription and translation leading to the production of a gene product. A gene product comprises either RNA or protein and is the result of expression of a gene. The amount of a gene product can be used to measure how active a gene is. The term "gene" as used in the present specification and in the claims comprises both coding regions (exons) as well as non-coding regions (e.g. non-coding regulatory elements such as promoters or enhancers, introns, leader and trailer sequences). The term "fragment" as used herein is meant to comprise e.g. an alternatively spliced, or truncated, or otherwise cleaved transcription product or translation product. The term "derivative" as used herein refers to a mutant, or an RNA-edited, or a chemically modified, or otherwise altered transcription product, or to a mutant, or chemically modified, or otherwise altered translation product. For instance, a "derivative" may be generated by processes such as altered phosphorylation, or glycosylation, or lipidation, or by altered signal peptide cleavage or other types of maturation cleavage. These processes may occur post-translationally. The term "modulator" as

used in the present invention and in the claims refers to a molecule capable of changing or altering the level and/or the activity of a gene, or a transcription product of a gene, or a translation product of a gene. Preferably, a "modulator" is capable of changing or altering the biological activity of a transcription product or a translation product of a gene. Said modulation, for instance, may be an increase or a decrease in enzyme activity, a change in binding characteristics, or any other change or alteration in the biological, functional, or immunological properties of said translation product of a gene. The term 'AD' shall mean Alzheimer's disease.

Neurodegenerative diseases or disorders according to the present invention comprise Alzheimer's disease, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, Pick's disease, fronto-temporal dementia, progressive nuclear palsy, corticobasal degeneration, cerebro-vascular dementia, multiple system atrophy, and mild-cognitive impairment. Further conditions involving neurodegenerative processes are, for instance, ischemic stroke, age-related macular degeneration and narcolepsy.

In one aspect, the invention features a method of diagnosing or prognosticating a neurodegenerative disease in a subject, or determining whether a subject is at increased risk of developing said disease. The method comprises: determining a level, or an activity, or both said level and said activity of (i) a transcription product of a gene coding for golgin-245, and/or of (ii) a translation product of a gene coding for golgin-245, and/or of (iii) a fragment or derivative of said transcription or translation product in a sample from said subject and comparing said level, and/or said activity to a reference value representing a known disease or health status, thereby diagnosing or prognosticating said neurodegenerative disease in said subject, or determining whether said subject is at increased risk of developing said neurodegenerative disease.

In a further aspect, the invention features a method of monitoring the progression of a neurodegenerative disease in a subject. A level, or an

activity, or both said level and said activity, of (i) a transcription product of a gene coding for golgin-245, and/or of (ii) a translation product of a gene coding for golgin-245, and/or of (iii) a fragment or derivative of said transcription or translation product in a sample from said subject is determined. Said level and/or said activity is compared to a reference value representing a known disease or health status. Thereby the progression of said neurodegenerative disease in said subject is monitored.

In still a further aspect, the invention features a method of evaluating a treatment for a neurodegenerative disease, comprising determining a level, or an activity, or both said level and said activity of (i) a transcription product of a gene coding for golgin-245, and/or of (ii) a translation product of a gene coding for a golgin-245, and/or of (iii) a fragment or derivative of said transcription or translation product in a sample obtained from a subject being treated for said disease. Said level, or said activity, or both said level and said activity are compared to a reference value representing a known disease or health status, thereby evaluating the treatment for said neurodegenerative disease.

In a preferred embodiment, said subjects suffer from Alzheimer's disease.

The present invention discloses the differential expression and regulation of the golgin-245 gene in specific brain regions of AD patients. Consequently, the golgin-245 gene and its corresponding transcription and translation products may have a causative role in the regional selective neuronal degeneration typically observed in AD. Alternatively, golgin-245 may confer a neuroprotective function to the remaining surviving nerve cells. Based on these disclosures, the present invention has utility for the diagnostic evaluation and prognosis as well as for the identification of a predisposition to a neurodegenerative disease, in particular AD. Furthermore, the present invention provides methods for the diagnostic monitoring of patients undergoing treatment for such a disease.

It is particularly preferred that said sample to be analyzed and determined is selected from the group consisting of a brain tissue or other tissues, organs or body cells. The sample can also consist of cerebrospinal fluid or other body fluids including saliva, urine, blood, serum plasma, or nasal mucosa.

In further preferred embodiments, said reference value is that of a level, or an activity, or both said level and said activity of (i) a transcription product of a gene coding for golgin-245, and/or of (ii) a translation product of a gene coding for golgin-245, and/or of (iii) a fragment or derivative of said transcription or translation product in a sample from a subject not suffering from said neurodegenerative disease.

In preferred embodiments, an increase or decrease in a transcription product of the gene coding for golgin-245 and /or a translation product of the gene coding for golgin-245 in a sample cell, or tissue, or body fluid from said subject relative to a reference value representing a known health status indicates a diagnosis, or prognosis, or increased risk of becoming diseased with a neurodegenerative disease, particularly AD.

In preferred embodiments, measurement of the level of transcription products of a gene coding for golgin-245 is performed in a sample from a subject using a quantitative PCR-analysis with primer combinations to amplify said gene specific sequences from cDNA obtained by reverse transcription of RNA extracted from a sample of a subject. A Northern blot with probes specific for said gene can also be applied. It might further be preferred to measure transcription products by means of chip-based micro-array technologies. These techniques are known to those of ordinary skill in the art (see Sambrook and Russell, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 2000).

Furthermore, a level and/or activity of a translation product of a gene coding for golgin-245 and/or fragment of said translation product can be detected using an immunoassay, an activity assay, and/or binding assay. These assays

can measure the amount of binding between said protein molecule and an anti-protein antibody by the use of enzymatic, chromodynamic, radioactive, magnetic, or luminescent labels which are attached to either the anti-protein antibody or a secondary antibody which binds the anti-protein antibody. In addition, other high affinity ligands may be used. Immunoassays which can be used include e.g. ELISAs, Western blots and other techniques known to those of ordinary skill in the art (see Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1999). All these detection techniques may also be employed in the format of micro-arrays, protein-arrays, or protein-chip based technologies.

In a preferred embodiment, the level, or the activity, or both said level and said activity of (i) a transcription product of a gene coding for golgin-245, and/or of (ii) a translation product of a gene coding for golgin-245, and/or of (iii) a fragment or derivative of said transcription or translation product in a series of samples taken from said subject over a period of time is compared, in order to monitor the progression of said disease. In further preferred embodiments, said subject receives a treatment prior to one or more of said sample gatherings. In yet another preferred embodiment, said level and/or activity is determined before and after said treatment of said subject.

In another aspect, the invention features a kit for diagnosing or prognosticating neurodegenerative diseases, in particular AD, in a subject, or determining the propensity or predisposition of a subject to develop a neurodegenerative disease, in particular AD, said kit comprising:

- (a) at least one reagent which is selected from the group consisting of (i) reagents that selectively detect a transcription product of a gene coding for golgin-245 (ii) reagents that selectively detect a translation product of a gene coding for golgin-245; and
- (b) instruction for diagnosing, or prognosticating a neurodegenerative disease, in particular AD, or determining the propensity or predisposition of a subject to develop such a disease by

- detecting a level, or an activity, or both said level and said activity, of said transcription product and/or said translation product of a gene coding for golgin-245, in a sample from said subject; and
- diagnosing or prognosticating a neurodegenerative disease, in particular AD, or determining the propensity or predisposition of said subject to develop such a disease,

wherein a varied level, or activity, or both said level and said activity, of said transcription product and/or said translation product compared to a reference value representing a known health status; or a level, or activity, or both said level and said activity, of said transcription product and/or said translation product similar or equal to a reference value representing a known disease status, indicates a diagnosis or prognosis of a neurodegenerative disease, in particular AD, or an increased propensity or predisposition of developing such a disease. The kit, according to the present invention, may be particularly useful for the identification of individuals that are at risk of developing a neurodegenerative disease, in particular AD. Consequently, the kit, according to the invention, may serve as a means for targeting identified individuals for early preventive measures or therapeutic intervention prior to disease onset, before irreversible damage in the course of the disease has been inflicted. Furthermore, in preferred embodiments, the kit featured in the invention is useful for monitoring a progression of a neurodegenerative disease, in particular AD in a subject, as well as monitoring success or failure of therapeutic treatment for such a disease of said subject.

In another aspect, the invention features a method of treating or preventing a neurodegenerative disease, in particular AD, in a subject comprising the administration to said subject in a therapeutically or prophylactically effective amount of an agent or agents which directly or indirectly affect a level, or an activity, or both said level and said activity, of (i) a gene coding for golgin-245, and/or (ii) a transcription product of a gene coding for golgin-245, and/or (iii) a translation product of a gene coding for golgin-245, and/or (iv) a fragment or derivative of (i) to (iii). Said agent may comprise a small molecule, or it may also comprise a peptide, an oligopeptide, or a

polypeptide. Said peptide, oligopeptide, or polypeptide may comprise an amino acid sequence of a translation product of a gene coding for golgin-245, or a fragment, or derivative, or a variant thereof.

In preferred embodiments, the method comprises the application of per se known methods of gene therapy and/or antisense nucleic acid technology to administer said agent or agents. In general, gene therapy includes several approaches: molecular replacement of a mutated gene, addition of a new gene resulting in the synthesis of a therapeutic protein, and modulation of endogenous cellular gene expression by recombinant expression methods or by drugs. Gene-transfer techniques are described in detail (see e.g. Behr, *Acc Chem Res* 1993, 26: 274-278 and Mulligan, *Science* 1993, 260: 926-931) and include direct gene-transfer techniques such as mechanical microinjection of DNA into a cell as well as indirect techniques employing biological vectors (like recombinant viruses, especially retroviruses) or model liposomes, or techniques based on transfection with DNA coprecipitation with polycations, cell membrane perturbation by chemical (solvents, detergents, polymers, enzymes) or physical means (mechanic, osmotic, thermic, electric shocks). The postnatal gene transfer into the central nervous system has been described in detail (see e.g. Wolff, *Curr Opin Neurobiol* 1993, 3: 743-748).

In particular, the invention features a method of treating or preventing a neurodegenerative disease by means of antisense nucleic acid therapy, i.e. the down-regulation of an inappropriately expressed or defective gene by the introduction of antisense nucleic acids or derivatives thereof into certain critical cells (see e.g. Gillespie, *DN&P* 1992, 5: 389-395; Agrawal and Akhtar, *Trends Biotechnol* 1995, 13: 197-199; Crooke, *Biotechnology* 1992, 10: 882-6). Apart from hybridization strategies, the application of ribozymes, i.e. RNA molecules that act as enzymes, destroying RNA that carries the message of disease has also been described (see e.g. Barinaga, *Science* 1993, 262: 1512-1514). In preferred embodiments, the subject to be treated is a human, and therapeutic antisense nucleic acids or derivatives thereof are directed against transcripts of a gene coding for golgin-245. It is preferred that cells of

the central nervous system, preferably the brain, of a subject are treated in such a way. Cell penetration can be performed by known strategies such as coupling of antisense nucleic acids and derivatives thereof to carrier particles, or the above described techniques. Strategies for administering targeted therapeutic oligo-deoxynucleotides are known to those of skill in the art (see e.g. Wickstrom, *Trends Biotechnol* 1992, 10: 281-287). In some cases, delivery can be performed by mere topical application. Further approaches are directed to intracellular expression of antisense RNA. In this strategy, cells are transformed *ex vivo* with a recombinant gene that directs the synthesis of an RNA that is complementary to a region of target nucleic acid. Therapeutical use of intracellularly expressed antisense RNA is procedurally similar to gene therapy.

In further preferred embodiments, the method comprises grafting donor cells into the central nervous system, preferably the brain, of said subject, or donor cells preferably treated so as to minimize or reduce graft rejection, wherein said donor cells are genetically modified by insertion of at least one transgene encoding said agent or agents. Said transgene might be carried by a viral vector, in particular a retroviral vector. The transgene can be inserted into the donor cells by a nonviral physical transfection of DNA encoding a transgene, in particular by microinjection. Insertion of the transgene can also be performed by electroporation, chemically mediated transfection, in particular calcium phosphate transfection, liposomal mediated transfection.

In preferred embodiments, said agent for treating and preventing a neurodegenerative disease, in particular AD, is a therapeutic protein which can be administered to said subject, preferably a human, by a process comprising introducing subject cells into said subject, said subject cells having been treated *in vitro* to insert a DNA segment encoding said therapeutic protein, said subject cells expressing *in vivo* in said subject a therapeutically effective amount of said therapeutic protein. Said DNA segment can be inserted into said cells *in vitro* by a viral vector, in particular a retroviral vector.

Methods of treatment, according to the present invention, comprise the application of therapeutic cloning, transplantation, and stem cell therapy using embryonic stem cells or embryonic germ cells and neuronal adult stem cells, combined with any of the previously described cell- and gene therapeutic methods. Stem cells may be totipotent or pluripotent. They may also be organ-specific. Strategies for repairing diseased and/or damaged brain cells or tissue comprise (i) taking donor cells from an adult tissue. Nuclei of those cells are transplanted into unfertilized egg cells from which the genetic material has been removed. Embryonic stem cells are isolated from the blastocyst stage of the cells which underwent somatic cell nuclear transfer. Use of differentiation factors then leads to a directed development of the stem cells to specialized cell types, preferably neuronal cells (Lanza et al., *Nature Medicine* 1999, 9: 975-977), or (ii) purifying adult stem cells, isolated from the central nervous system, or from bone marrow (mesenchymal stem cells), for in vitro expansion and subsequent grafting and transplantation, or (iii) directly inducing endogenous neural stem cells to proliferate, migrate, and differentiate into functional neurons (Peterson DA, *Curr. Opin. Pharmacol.* 2002, 2: 34-42). Adult neural stem cells are of great potential for repairing damaged or diseased brain tissues, as the germinal centers of the adult brain are free of neuronal damage or dysfunction (Colman A, *Drug Discovery World* 2001, 7: 66-71).

In preferred embodiments, the subject for treatment or prevention, according to the present invention, can be a human, an experimental animal, e.g. a mouse or a rat, a domestic animal, or a non-human primate. The experimental animal can be an animal model for a neurodegenerative disorder, e.g. a transgenic mouse and/or a knock-out mouse with an AD-type neuropathology.

In a further aspect, the invention features a modulator of an activity, or a level, or both said activity and said level of at least one substance which is selected from the group consisting of (i) a gene coding for golgin-245; and/or (ii) a transcription product of a gene coding for a golgin-245, and/or (iii) a

translation product of a gene coding for golgin-245, and/or (iv) a fragment or derivative of (i) to (iii).

In an additional aspect, the invention features a pharmaceutical composition comprising said modulator and preferably a pharmaceutical carrier. Said carrier refers to a diluent, adjuvant, excipient, or vehicle with which the modulator is administered.

In a further aspect, the invention features a modulator of an activity, or a level, or both said activity and said level of at least one substance which is selected from the group consisting of (i) a gene coding for golgin-245, and/or (ii) a transcription product of a gene coding for golgin-245, and/or (iii) a translation product of a gene coding for golgin-245, and/or (iv) a fragment or derivative of (i) to (iii) for use in a pharmaceutical composition.

In another aspect, the invention provides for the use of a modulator of an activity, or a level, or both said activity and said level of at least one substance which is selected from the group consisting of (i) a gene coding for golgin-245, and/or (ii) a transcription product of a gene coding for golgin-245 and/or (iii) a translation product of a gene coding for golgin-245, and/or (iv) a fragment or derivative of (i) to (iii) for a preparation of a medicament for treating or preventing a neurodegenerative disease, in particular AD.

In one aspect, the present invention also provides a kit comprising one or more containers filled with a therapeutically or prophylactically effective amount of said pharmaceutical composition.

In a further aspect, the invention features a recombinant, non-human animal comprising a non-native gene sequence coding for golgin-245, or a fragment thereof, or a derivative thereof. The generation of said recombinant, non-human animal comprises (i) providing a gene targeting construct containing said gene sequence and a selectable marker sequence, and (ii) introducing said targeting construct into a stem cell of a non-human animal, and (iii)

introducing said non-human animal stem cell into a non-human embryo, and (iv) transplanting said embryo into a pseudopregnant non-human animal, and (v) allowing said embryo to develop to term, and (vi) identifying a genetically altered non-human animal whose genome comprises a modification of said gene sequence in both alleles, and (vii) breeding the genetically altered non-human animal of step (vi) to obtain a genetically altered non-human animal whose genome comprises a modification of said endogenous gene, wherein said gene is mis-expressed, or under-expressed, or over-expressed, and wherein said disruption or alteration results in said non-human animal exhibiting a predisposition to developing symptoms of neuropathology similar to a neurodegenerative disease, in particular AD. Strategies and techniques for the generation and construction of such an animal are known to those of ordinary skill in the art (see e.g. Capecchi, *Science* 1989, 244: 1288-1292 and Hogan et al., 1994, *Manipulating the Mouse Embryo: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York). It is preferred to make use of such a recombinant non-human animal as an animal model for investigating neurodegenerative diseases, in particular AD.

In another aspect, the invention features an assay for screening for a modulator of neurodegenerative diseases, in particular AD, or related diseases and disorders of one or more substances selected from the group consisting of (i) a gene coding for golgin-245, and/or (ii) a transcription product of a gene coding for golgin-245, and/or (iii) a translation product of a gene coding for golgin-245, and/or (iv) a fragment or derivative of (i) to (iii). This screening method comprises (a) contacting a cell with a test compound, and (b) measuring the activity, or the level, or both the activity and the level of one or more substances recited in (i) to (iv), and (c) measuring the activity, or the level, or both the activity and the level of said substances in a control cell not contacted with said test compound, and (d) comparing the levels of the substance in the cells of step (b) and (c), wherein an alteration in the activity and/or level of said substances in the contacted cells indicates that the test compound is a modulator of said diseases and disorders.

In one further aspect, the invention features a screening assay for a modulator of neurodegenerative diseases, in particular AD, or related diseases and disorders of one or more substances selected from the group consisting of (i) a gene coding for golgin-245, and/or (ii) a transcription product of a gene coding for golgin-245, and/or (iii) a translation product of a gene coding for golgin-245, and/or (iv) a fragment or derivative of (i) to (iii), comprising (a) administering a test compound to a test animal which is predisposed to developing or has already developed a neurodegenerative disease or related diseases or disorders, and (b) measuring the activity and/or level of one or more substances recited in (i) to (iv), and (c) measuring the activity and/or level of said substances in a matched control animal which is equally predisposed to developing or has already developed said diseases and to which animal no such test compound has been administered, and (d) comparing the activity and/or level of the substance in the animals of step (b) and (c), wherein an alteration in the activity and/or level of substances in the test animal indicates that the test compound is a modulator of said diseases and disorders.

In a preferred embodiment, said test animal and/or said control animal is a recombinant, non-human animal which expresses a gene coding for golgin-245, or a fragment thereof, or a derivative thereof, under the control of a transcriptional regulatory element which is not the native golgin-245 gene transcriptional control regulatory element.

In another embodiment, the present invention provides a method for producing a medicament comprising the steps of (i) identifying a modulator of neurodegenerative diseases by a method of the aforementioned screening assays and (ii) admixing the modulator with a pharmaceutical carrier. However, said modulator may also be identifiable by other types of screening assays.

In another aspect, the present invention provides for an assay for testing a compound, preferably for screening a plurality of compounds, for inhibition of

binding between a ligand and golgin-245 protein, or a fragment or derivative thereof. Said screening assay comprises the steps of (i) adding a liquid suspension of said golgin-245 protein, or a fragment or derivative thereof, to a plurality of containers, and (ii) adding a compound or a plurality of compounds to be screened for said inhibition to said plurality of containers, and (iii) adding fluorescently labelled ligand to said containers, and (iv) incubating said golgin-245 protein, or said fragment or derivative thereof, and said compound or plurality of compounds, and said fluorescently labelled ligand, and (v) measuring the amounts of fluorescence associated with said golgin-245 protein, or with said fragment or derivative thereof, and (vi) determining the degree of inhibition by one or more of said compounds of binding of said ligand to said golgin-245 protein, or said fragment or derivative thereof. Instead of utilizing a fluorescently labelled ligand, it might in some aspects be preferred to use any other detectable label known to the person skilled in the art, e.g. radioactive labels, and detect it accordingly. Said method may be useful for the identification of novel compounds as well as for evaluating compounds which have been improved or otherwise optimized in their ability to inhibit the binding of a ligand to a gene product of a gene coding for golgin-245, or a fragment or derivative thereof.

In one further embodiment, the present invention provides a method for producing a medicament comprising the steps of (i) identifying a compound as an inhibitor of binding between a ligand and a gene product of a gene coding for golgin-245 by the aforementioned inhibitory binding assay and (ii) admixing the compound with a pharmaceutical carrier. However, said compound may also be identifiable by other types of screening assays.

In another aspect, the invention features an assay for testing a compound, preferably for screening a plurality of compounds to determine the degree of binding of said compounds to golgin-245 protein, or to a fragment or derivative thereof. Said screening assay comprises (i) adding a liquid suspension of said golgin-245 protein, or a fragment or derivative thereof, to a plurality of containers, and (ii) adding a fluorescently labelled compound or a

plurality of fluorescently labelled compounds to be screened for said binding to said plurality of containers, and (iii) incubating said golgin-245 protein, or said fragment or derivative thereof, and said fluorescently labelled compound or fluorescently labelled compounds, and (iv) measuring the amounts of fluorescence associated with said golgin-245 protein, or with said fragment or derivative thereof, and (v) determining the degree of binding by one or more of said compounds to said golgin-245 protein, or said fragment or derivative thereof. In this type of assay it might be preferred to use a fluorescent label. However, any other type of detectable label might also be employed. Said method may be useful for the identification of novel compounds as well as for evaluating compounds which have been improved or otherwise optimized in their ability to bind to golgin-245, or a fragment or derivative thereof.

In one further embodiment, the present invention provides a method for producing a medicament comprising the steps of (i) identifying a compound as a binder to a gene product of a gene coding for golgin-245 by the aforementioned binding assays and (ii) admixing the compound with a pharmaceutical carrier. However, said compound may also be identifiable by other types of screening assays.

In another embodiment, the present invention provides for a medicament obtainable by any of the methods according to the herein claimed screening assays. In one further embodiment, the instant invention provides for a medicament obtained by any of the methods according to the herein claimed screening assays.

The present invention features an antibody which is specifically immunoreactive with an immunogen, wherein said immunogen is a translation product of a gene coding for golgin-245, or a fragment or derivative thereof. The immunogen may comprise immunogenic or antigenic epitopes or portions of a translation product of said gene, wherein said immunogenic or antigenic portion of a translation product is a polypeptide, and wherein said polypeptide elicits an antibody response in an animal, and wherein said polypeptide is

Immunospecifically bound by said antibody. Methods for generating antibodies are well known in the art (see Harlow et al., *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York). The term "antibody", as employed in the present invention, encompasses all forms of antibodies known in the art, such as polyclonal, monoclonal, chimeric, recombinatorial, anti-idiotypic, humanized, or single chain antibodies, as well as fragments thereof. Antibodies of the present invention are useful, for instance, in a variety of diagnostic and therapeutic methods involving detecting translation products of a gene coding for golgin-245.

In a preferred embodiment of the present invention, said antibodies can be used for detecting the pathological state of a cell in a sample from a subject, comprising immunocytochemical staining of said cell with said antibody, wherein an altered degree of staining, or an altered staining pattern in said cell compared to a cell representing a known health status indicates a pathological state of said cell. Preferably, the pathological state relates to a neurodegenerative disease, in particular to AD. Immunocytochemical staining of a cell can be carried out by a number of different experimental methods well known in the art. It might be preferred, however, to apply an automated method for the detection of antibody binding, wherein the determination of the degree of staining of a cell, or the determination of the cellular or subcellular staining pattern of a cell, or the topological distribution of an antigen on the cell surface or among organelles and other subcellular structures within the cell, are carried out according to the method described in US patent 6150173).

Other features and advantages of the invention will be apparent from the following description of figures and examples which are illustrative only and not intended to limit the remainder of the disclosure in any way.

Figure 1 depicts the brain regions with selective vulnerability to neuronal loss and degeneration in AD. Primarily, neurons within the inferior temporal lobe;

the entorhinal cortex, the hippocampus, and the amygdala are subject to degenerative processes in AD (Terry et al., *Annals of Neurology* 1981, 10:184-192). These brain regions are mostly involved in the processing of learning and memory functions. In contrast, neurons within the frontal cortex, the occipital cortex, and the cerebellum remain largely intact and preserved from neurodegenerative processes in AD. Brain tissues from the frontal cortex (F) and the temporal cortex (T) of AD patients and healthy, age-matched control individuals were used for the herein disclosed examples. For illustrative purposes, the image of a normal healthy brain was taken from a publication by Strange (*Brain Biochemistry and Brain Disorders*, Oxford University Press, Oxford, 1992, p.4).

Figure 2 discloses the initial identification of the differential expression of the gene coding for golgin-245 in a fluorescence differential display screen. The figure shows a clipping of a large preparative fluorescent differential display gel. PCR products from the frontal cortex (F) and the temporal cortex (T) of two healthy control subjects and six AD patients were loaded in duplicate onto a denaturing polyacrylamide gel (from left to right). PCR products were obtained by amplification of the individual cDNAs with the corresponding one-base-anchor oligonucleotide and the specific Cy3 labelled random primers. The arrow indicates the migration position where significant differences in intensity of the signals for a transcription product of the gene coding for golgin-245 derived from frontal cortex as compared to the signals derived from the temporal cortex of AD patients exist. The differential expression reflects an up-regulation of golgin-245 gene transcription in the temporal cortex compared to the frontal cortex of AD patients. Comparing the signals derived from temporal cortex and frontal cortex of healthy non-AD control subjects with each other, no difference in signal intensity, i.e. no altered expression level can be detected.

Figure 3 illustrates the verification of the differential expression of the gene coding for golgin-245 by quantitative RT-PCR analysis. Quantification of RT-PCR products from RNA samples collected from the frontal cortex (F) and

temporal cortex (T) of AD patients (Fig 3b) and of healthy, age-matched control individuals (Fig 3a) was performed by the LightCycler rapid thermal cycling technique. The data were normalized to the combined average values of a set of standard genes which showed no significant differences in their gene expression levels. Said set of standard genes consisted of genes for the ribosomal protein S9, the transferrin receptor, GAPDH, and beta-actin. The figure depicts the kinetics of amplification by plotting the cycle number against the amount of amplified material as measured by its fluorescence. Note that the amplification kinetics of the golgin-245 cDNA from both the frontal and temporal cortices of a normal control individual during the exponential phase of the reaction overlap (Fig 3a, arrow), whereas in AD (Fig 3b, arrows), there is a significant shift of the curve for the sample derived from temporal cortex, indicating an up-regulation of a transcription product of the gene coding for golgin-245 in temporal cortex relative to frontal cortex.

Figure 4 discloses SEQ ID NO. 1, the polypeptide sequence of golgin-245 comprising 2230 amino acids. The protein harbors several distinct functional domains which are located as follows: amino acid residues 1 to 139 and 261 to 292 form proline-rich domains, amino acid residues 555 to 564 generate the granine signature, and the Golgi-targeting signal spans amino acids 2180-2250 containing the highly conserved tyrosine residue Y2199.

Figure 5 represents SEQ ID NO. 2, the nucleotide sequence of golgin-245 cDNA, comprising 7743 nucleotides.

Figure 6 depicts SEQ ID NO. 3, the nucleotide sequence of the 36 bp golgin-245 cDNA fragment, identified and obtained by fluorescence differential display and subsequent cloning.

Figure 7 outlines the sequence alignment of SEQ ID NO. 3, the 36 bp human golgin-245 cDNA fragment, with SEQ ID NO. 2, the nucleotide sequence of golgin-245 cDNA (nucleotides 5554 to 5589).

Table 1 lists the gene expression levels in the temporal cortex relative to the frontal cortex for the golgin-245 gene in seven Alzheimer's disease patients (0.98-2.91 Δ -fold) and four healthy, age-matched control individuals (0.86-1.00 Δ -fold).

EXAMPLE I:

(i) Brain tissue dissection from patients with AD:

Brain tissues from AD patients and age-matched control subjects were collected within 6 hours post-mortem and immediately frozen on dry ice. Sample sections from each tissue were fixed in paraformaldehyde for histopathological confirmation of the diagnosis. Brain areas for differential expression analysis were identified (see Fig. 1) and stored at -80°C until RNA extractions were performed.

(ii) Isolation of total mRNA:

Total RNA was extracted from post-mortem brain tissue by using the RNeasy kit (Qiagen) according to the manufacturer's protocol. The accurate RNA concentration and the RNA quality were determined with the DNA LabChip system using the Agilent 2100 Bioanalyzer (Agilent Technologies). For additional quality testing of the prepared RNA, i.e. exclusion of partial degradation and testing for DNA contamination, specifically designed intronic GAPDH oligonucleotides and genomic DNA as reference control were utilised to generate a melting curve with the LightCycler technology as described in the supplied protocol by the manufacturer (Roche).

(iii) cDNA synthesis and identification of differentially expressed genes by fluorescence differential display (FDD):

In order to identify changes in gene expression in different tissues we employed a modified and improved differential display (DD) screening method. The original DD screening method is known to those skilled in the art (Liang and Pardee, *Science* 1995, 267:1186-7). This technique compares two populations of RNA and provides clones of genes that are expressed in one

population but not in the other. Several samples can be analyzed simultaneously and both up- and down-regulated genes can be identified in the same experiment. By adjusting and refining several steps in the DD method as well as modifying technical parameters, e.g. increasing redundancy, evaluating optimized reagents and conditions for reverse transcription of total RNA, optimizing polymerase chain reactions (PCR) and separation of the products thereof, a technique was developed which allows for highly reproducible and sensitive results. The applied and improved DD technique was described in detail by von der Kammer et al. (*Nucleic Acids Research* 1999, 27: 2211-2218). A set of 64 specifically designed random primers was developed (standard set) to achieve a statistically comprehensive analysis of all possible RNA species. Further, the method was modified to generate a preparative DD slab-gel technique, based on the use of fluorescently labelled primers. In the present invention, RNA populations from carefully selected post-mortem brain tissues (frontal and temporal cortex) of AD patients and age-matched control subjects were compared.

As starting material for the DD analysis we used total RNA, extracted as described above (ii). Equal amounts of 0.05 µg RNA each were transcribed into cDNA in 20 µl reactions containing 0.5 mM each dNTP, 1 µl Sensiscript Reverse Transcriptase and 1x RT buffer (Qiagen), 10 U RNase inhibitor (Qiagen) and 1 µM of either one-base-anchor oligonucleotides HT11A, HT11G or HT11C (Liang et al., *Nucleic Acids Research* 1994, 22: 5763-5764; Zhao et al., *Biotechniques* 1995, 18: 842-850). Reverse transcription was performed for 60 min at 37 °C with a final denaturation step at 93 °C for 5 min: 2 µl of the obtained cDNA each was subjected to a polymerase chain reaction (PCR) employing the corresponding one-base-anchor oligonucleotide (1 µM) along with either one of the Cy3 labelled random DD primers (1 µM), 1x GeneAmp PCR buffer (Applied Biosystems), 1.5 mM MgCl₂ (Applied Biosystems), 2 µM dNTP-Mix (dATP, dGTP, dCTP, dTTP Amersham Pharmacia Biotech), 5 % DMSO (Sigma), 1 U AmpliTaq DNA Polymerase (Applied Biosystems) in a 20 µl final volume. PCR conditions were set as follows: one round at 94 °C for 30 sec for denaturing, cooling 1 °C/sec down to 40 °C, 40 °C for 4 min for low-

stringency annealing of primer, heating 1 °C/sec up to 72 °C, 72 °C for 1 min for extension. This round was followed by 39 high-stringency cycles: 94 °C for 30 sec, cooling 1 °C/sec down to 60 °C, 60 °C for 2 min, heating 1 °C/sec up to 72 °C, 72 °C for 1 min. One final step at 72 °C for 5 min was added to the last cycle (PCR cycler: Multi Cycler PTC 200, MJ Research). 8 µl DNA loading buffer were added to the 20 µl PCR product preparation, denatured for 5 min and kept on ice until loading onto a gel. 3.5 µl each were separated on 0.4 mm thick, 6 %-polyacrylamide (Long Ranger)/ 7 M urea sequencing gels in a slab-gel system (Hitachi Genetic Systems) at 2000 V, 60W, 30 mA, for 1 h 40 min. Following completion of the electrophoresis, gels were scanned with a FMBIO II fluorescence-scanner (Hitachi Genetic Systems), using the appropriate FMBIO II Analysis 8.0 software. A full-scale picture was printed, differentially expressed bands marked, excised from the gel, transferred into 1.5 ml containers, overlaid with 200 µl sterile water and kept at -20°C until extraction.

Elution and reamplification of DD products: The differential bands were extracted from the gel by boiling in 200 µl H₂O for 10 min, cooling down on ice and precipitation from the supernatant fluids by using ethanol (Merck) and glycogen/sodium acetate (Merck) at - 20 °C over night, and subsequent centrifugation at 13.000 rpm for 25 min at 4 °C. Pellets were washed twice in ice-cold ethanol (80%), resuspended in 10 mM Tris pH 8.3 (Merck) and dialysed against 10 % glycerol (Merck) for 1 h at room temperature on a 0.025 µm VSWP membrane (Millipore). The obtained preparations were used as templates for reamplification by 15 high-stringency cycles in 25-µl PCR mixtures containing the corresponding primer pairs as used for the DD PCR (see above) under identical conditions, with the exception of the initial round at 94 °C for 5 min, followed by 15 cycles of: 94 °C for 45 sec, 60 °C for 45 sec, ramp 1°C/sec to 70 °C for 45 sec, and one final step at 72 °C for 5 min.

Cloning and sequencing of DD products: Re-amplified cDNAs were analyzed with the DNA LabChip® system (Agilent 2100 Bioanalyzer, Agilent Technologies) and ligated into the pCR-Blunt II-TOPO vector and transformed into *E.coli* Top10F' cells (Zero Blunt TOPO PCR Cloning Kit, Invitrogen) according to the manufacturer's instructions. Cloned cDNA fragments were

sequenced by commercially available sequencing facilities. The result of one such FDD experiment for the golgin-245 gene is shown in Fig. 2.

(iv) Confirmation of differential expression by quantitative RT-PCR:

Positive corroboration of differential expression of the golgin-245 gene was performed using the LightCycler technology (Roche). This technique features rapid thermal cycling for the polymerase chain reaction as well as real-time measurement of fluorescent signals during amplification and therefore allows for highly accurate quantification of RT-PCR products by using a kinetic, rather than an endpoint readout. The ratio of golgin-245 cDNA from the temporal cortex relative to frontal cortex was determined (relative quantification).

First, a standard curve was generated to determine the efficiency of the PCR with specific primers for golgin-245 gene (5'-AGATGCTCGGCTGATGT-3' and 5'-GACATTGGGTGACTGCTGCTT-3'). PCR amplification (95 °C and 1 sec, 56 °C and 5 sec, and 72 °C and 5 sec) was performed in a volume of 20 µl containing LightCycler-FastStart DNA Master SYBR Green I mix (contains FastStart Taq DNA polymerase, reaction buffer, dNTP mix with dUTP instead of dTTP, SYBR Green I dye, and 1 mM MgCl₂; Roche), 0.5 µM primers, 2 µl of a cDNA dilution series (final concentration of 40, 20, 10, 5, 1 and 0.5 ng human total brain cDNA; Clontech) and, depending on the primers used, additional 3 mM MgCl₂. Melting curve analysis revealed a single peak at approximately 82°C with no visible primer dimers. Quality and size of the PCR product were determined with the DNA LabChip system (Agilent 2100 Bioanalyzer, Agilent Technologies). A single peak at the expected size of 132 bp for the golgin-245 gene was observed in the electropherogram of the sample.

In an analogous manner, the PCR protocol was applied to determine the PCR efficiency of a set of reference genes which were selected as a reference standard for quantification. In the present invention, the mean value of five such reference genes was determined: (1) cyclophilin B, using the specific primers 5'-ACTGAAGCACTACGGGCCTG-3' and 5'-AGCCGTTGGTGTCTT-TGCC-3' except for MgCl₂ (an additional 1 mM was added instead of 3 mM).

Melting curve analysis revealed a single peak at approximately 87 °C with no visible primer dimers. Agarose gel analysis of the PCR product showed one single band of the expected size (62 bp). (2) Ribosomal protein S9 (RPS9), using the specific primers 5'-GGTCAAATTTACCCTGGCCA-3' and 5'-TCTCATCAAGCGTCAGCAGTTC-3' (exception: additional 1 mM MgCl₂ was added instead of 3 mM). Melting curve analysis revealed a single peak at approximately 85°C with no visible primer dimers. Agarose gel analysis of the PCR product showed one single band with the expected size (62 bp). (3) beta-actin, using the specific primers 5'-TGGAACGGTGAAGGTGACA-3' and 5'-GGCAAGGGACTTCCTGTAA-3'. Melting curve analysis revealed a single peak at approximately 87°C with no visible primer dimers. Agarose gel analysis of the PCR product showed one single band with the expected size (142 bp). (4) GAPDH, using the specific primers 5'-CGTCATGGGTGTGAACCATG-3' and 5'-GCTAAGCAGTTGGTGGTGCAG-3'. Melting curve analysis revealed a single peak at approximately 83°C with no visible primer dimers. Agarose gel analysis of the PCR product showed one single band with the expected size (81 bp). (5) Transferrin receptor TRR, using the specific primers 5'-GTCGCTGGTCAGTTCGTGATT-3' and 5'-AGCAGTTGGCTGTTGTACCTCTC-3'. Melting curve analysis revealed a single peak at approximately 83°C with no visible primer dimers. Agarose gel analysis of the PCR product showed one single band with the expected size (80 bp).

For calculation of the values, first the logarithm of the cDNA concentration was plotted against the threshold cycle number C_t for golgin-245 and the five reference standard genes. The slopes and the intercepts of the standard curves (i.e. linear regressions) were calculated for all genes. In a second step, cDNAs from frontal cortex and temporal cortex were analyzed in parallel and normalized to cyclophilin B. The C_t values were measured and converted to ng total brain cDNA using the corresponding standard curves:

$$10^{((C_t \text{ value} - \text{intercept}) / \text{slope})} \quad [\text{ng total brain cDNA}]$$

The values for frontal and temporal cortex cDNAs of golgin-245 cDNA were normalized to cyclophilin B, and the ratio was calculated using the following formula:

$$\text{Ratio} = \frac{\text{golgin-245 temporal [ng]} / \text{cyclophilin B temporal [ng]}}{\text{golgin-245 frontal [ng]} / \text{cyclophilin B frontal [ng]}}$$

In a third step, the set of reference standard genes was analyzed in parallel to determine the mean average value of the temporal to frontal ratios of expression levels of the reference standard genes for each individual brain sample. As cyclophilin B was analyzed in step 2 and step 3, and the ratio from one gene to another gene remained constant in different runs, it was possible to normalize the values for golgin-245 to the mean average value of the set of reference standard genes instead of normalizing to one single gene alone. The calculation was performed by dividing the ratio shown above by the deviation of cyclophilin B from the mean value of all housekeeping genes. The results of one such quantitative RT-PCR analysis for the golgin-245 gene are shown in Figure 3.

CLAIMS

1. A method of diagnosing or prognosticating a neurodegenerative disease in a subject, or determining whether a subject is at increased risk of developing said disease, comprising:
determining a level and/or an activity of
 - (i) a transcription product of a gene coding for golgin-245, and/or
 - (ii) a translation product of a gene coding for golgin-245 and/or
 - (iii) a fragment or derivative of said transcription or translation productin a sample from said subject and comparing said level and/or said activity to a reference value representing a known disease or health status, thereby diagnosing or prognosticating said neurodegenerative disease in said subject, or determining whether said subject is at increased risk of developing said neurodegenerative disease.
2. A method of monitoring the progression of a neurodegenerative disease in a subject, comprising:
determining a level and/or an activity of
 - (i) a transcription product of a gene coding for golgin-245, and/or
 - (ii) a translation product of a gene coding for golgin-245, and/or
 - (iii) a fragment or derivative of said transcription or translation productin a sample from said subject and comparing said level and/or said activity to a reference value representing a known disease or health status, thereby monitoring the progression of said neurodegenerative disease in said subject.
3. A method of evaluating a treatment for a neurodegenerative disease, comprising:
determining a level and/or an activity of
 - (i) a transcription product of a gene coding for golgin-245, and/or
 - (ii) a translation product of a gene coding for golgin-245, and/or
 - (iii) a fragment or derivative of said transcription or translation productin a sample from a subject being treated for said disease and comparing said level and/or said activity to a reference value representing a known disease or

health status, thereby evaluating said treatment for said neurodegenerative disease.

4. The method according to any of claims 1 to 3 wherein said neurodegenerative disease is Alzheimer's disease.

5. The method according to any of claims 1 to 4 wherein said sample is a cell, or a tissue, or an organ, or a body fluid, in particular cerebrospinal fluid or blood.

6. The method according to any of claims 1 to 5 wherein said reference value is that of a level and/or an activity of

- (i) a transcription product of a gene coding for golgin-245, and/or
 - (ii) a translation product of a gene coding for golgin-245, and/or
 - (iii) a fragment or derivative of said transcription or translation product
- in a sample from a subject not suffering from said neurodegenerative disease.

7. The method according to any of claims 1 to 6 wherein an increase or decrease in a transcription product of the gene coding for golgin-245 and/or a translation product of a gene coding for golgin-245 in a cell, or tissue, or body fluid from said subject relative to a reference value representing a known health status indicates a diagnosis, or prognosis, or increased risk of Alzheimer's disease in said subject.

8. The method according to any of claims 1 to 7, further comprising comparing a level and/or an activity of

- (i) a transcription product of a gene coding for golgin-245, and/or
 - (ii) a translation product of a gene coding for golgin-245, and/or
 - (iii) a fragment or derivative of said transcription or translation product
- in a series of samples taken from said subject over a period of time.

9. The method according to claim 8 wherein said subject receives a treatment prior to one or more of said sample gatherings.

10. The method according to claim 9 wherein said level and/or activity is determined before and after said treatment of said subject.

11. A kit for diagnosing or prognosticating a neurodegenerative disease, in particular Alzheimer's disease, in a subject, or determining the propensity or predisposition of a subject to develop such a disease, said kit comprising:

- (a) at least one reagent which is selected from the group consisting of (i) reagents that selectively detect a transcription product of a gene coding for golgin-245 and (ii) reagents that selectively detect a translation product of a gene coding for golgin-245 and
- (b) an instruction for diagnosing, or prognosticating a neurodegenerative disease, in particular Alzheimer's disease, or determining the propensity or predisposition of a subject to develop such a disease by (i) detecting a level, or an activity, or both said level and said activity, of said transcription product and/or said translation product of a gene coding for golgin-245, in a sample from said subject; and (ii) diagnosing or prognosticating a neurodegenerative disease, in particular Alzheimer's disease, or determining the propensity or predisposition of said subject to develop such a disease, wherein a varied level, or activity, or both said level and said activity, of said transcription product and/or said translation product compared to a reference value representing a known health status; or a level, or activity, or both said level and said activity, of said transcription product and/or said translation product similar or equal to a reference value representing a known disease status indicates a diagnosis or prognosis of a neurodegenerative disease, in particular Alzheimer's disease, or an increased propensity or predisposition of developing such a disease.

12. A method of treating or preventing a neurodegenerative disease, in particular Alzheimer's disease, in a subject comprising administering to said subject in a therapeutically or prophylactically effective amount an agent or agents which directly or indirectly affect an activity and/or a level of

- (i) a gene coding for golgin-245, and/or
- (ii) a transcription product of a gene coding for golgin-245, and/or
- (iii) a translation product of a gene coding for golgin-245, and/or
- (iv) a fragment or derivative of (i) to (iii).

13. A modulator of an activity and/or of a level of at least one substance which is selected from the group consisting of

- (i) a gene coding for golgin-245 and/or
- (ii) a transcription product of a gene coding for golgin-245 and/or
- (iii) a translation product of a gene coding for golgin-245, and/or
- (iv) a fragment or derivative of (i) to (iii).

14. A recombinant, non-human animal comprising a non-native gene sequence coding for golgin-245 or a fragment thereof, or a derivative thereof, said animal being obtainable by:

- (i) providing a gene targeting construct comprising said gene sequence and a selectable marker sequence, and
- (ii) introducing said targeting construct into a stem cell of a non-human animal; and
- (iii) introducing said non-human animal stem cell into a non-human embryo, and
- (iv) transplanting said embryo into a pseudopregnant non-human animal, and
- (v) allowing said embryo to develop to term, and
- (vi) identifying a genetically altered non-human animal whose genome comprises a modification of said gene sequence in both alleles, and
- (vii) breeding the genetically altered non-human animal of step (vi) to obtain a genetically altered non-human animal whose genome comprises a modification of said endogenous gene, wherein said disruption results in said non-human animal exhibiting a predisposition to developing a neurodegenerative disease or related diseases or disorders.

15. An assay for screening for a modulator of neurodegenerative diseases, in particular Alzheimer's disease, or related diseases or disorders of one or more substances selected from the group consisting of

- (i) a gene coding for golgin-245, and/or
- (ii) a transcription product of a gene coding for golgin-245, and/or
- (iii) a translation product of a gene coding for golgin-245, and/or
- (iv) a fragment or derivative of (i) to (iii),

said method comprising:

- (a) contacting a cell with a test compound;
- (b) measuring the activity and/or level of one or more substances recited in (i) to (iv);
- (c) measuring the activity and/or level of one or more substances recited in (i) to (iv) in a control cell not contacted with said test compound; and
- (d) comparing the levels and/or activities of the substance in the cells of step (b) and (c), wherein an alteration in the activity and/or level of substances in the contacted cells indicates that the test compound is a modulator of said diseases or disorders.

16. A method of screening for a modulator of neurodegenerative diseases, in particular Alzheimer's disease, or related diseases or disorders of one or more substances selected from the group consisting of

- (i) a gene coding for golgin-245, and/or
- (ii) a transcription product of a gene coding for golgin-245, and/or
- (iii) a translation product of a gene coding for golgin-245, and/or
- (v) a fragment or derivative of (i) to (iii),

said method comprising:

- (a) administering a test compound to a test animal which is predisposed to developing or has already developed a neurodegenerative disease or related diseases or disorders in respect of the substances recited in (i) to (iv);
- (b) measuring the activity and/or level of one or more substances recited in (i) to (iv);

- (c) measuring the activity and/or level of one or more substances recited in (i) or (iv) in a matched control animal which is predisposed to developing or has already developed a neurodegenerative disease or related diseases or disorders in respect to the substances recited in (i) to (iv) and to which animal no such test compound has been administered;
- (d) comparing the activity and/or level of the substance in the animals of step (b) and (c), wherein an alteration in the activity and/or level of substances in the test animal indicates that the test compound is a modulator of said diseases or disorders.

17. The method according to claim 16 wherein said test animal and/or said control animal is a recombinant animal which expresses the gene coding for golgin-245, or a fragment thereof, or a derivative thereof, under the control of a transcriptional control element which is not the native golgin-245 gene transcriptional control element.

18. An assay for testing a compound, preferably for screening a plurality of compounds for inhibition of binding between a ligand and golgin-245 protein, or a fragment or derivative thereof, said assay comprising the steps of:

- (i) adding a liquid suspension of said golgin-245 protein, or a fragment or derivative thereof, to a plurality of containers;
- (ii) adding a compound or a plurality of compounds to be screened for said inhibition to said plurality of containers;
- (iii) adding fluorescently labelled ligand to said containers;
- (iv) incubating said golgin-245 protein, or said fragment or derivative thereof, and said compound or compounds, and said fluorescently labelled ligand;
- (v) measuring amounts of fluorescence associated with said golgin-245 protein, or with said fragment or derivative thereof; and
- (vi) determining the degree of inhibition by one or more of said compounds of binding of said ligand to said golgin-245 protein, or said fragment or derivative thereof.

19. An assay for testing a compound, preferably for screening a plurality of compounds to determine the degree of binding of said compounds to golgin-245 protein, or to a fragment or derivative thereof, said assay comprising the steps of:

- (i) adding a liquid suspension of said golgin-245 protein, or a fragment or derivative thereof, to a plurality of containers;
- (ii) adding a fluorescently labelled compound or a plurality of fluorescently labelled compounds to be screened for said binding to said plurality of containers;
- (iii) incubating said golgin-245 protein, or said fragment or derivative thereof, and said fluorescently labelled compound or fluorescently labelled compounds;
- (iv) measuring amounts of fluorescence associated with said golgin-245 protein, or with said fragment or derivative thereof; and
- (v) determining the degree of binding by one or more of said compounds to said golgin-245 protein, or said fragment or derivative thereof.


20. Use of an antibody specifically immunoreactive with an immunogen, wherein said immunogen is a translation product of a gene coding for golgin-245, or a fragment or derivative thereof, for detecting the pathological state of a cell in a sample from a subject; comprising immunocytochemical staining of said cell with said antibody, wherein an altered degree of staining, or an altered staining pattern in said cell compared to a cell representing a known health status indicates a pathological state of said cell, and wherein said pathological state relates to a neurodegenerative disease, in particular Alzheimer's disease.

SUMMARY

The present invention discloses the differential expression of the golgin-245 gene in specific brain regions of Alzheimer's disease patients. Based on this finding, this invention provides a method for diagnosing or prognosticating a neurodegenerative disease, in particular Alzheimer's disease, in a subject, or for determining whether a subject is at increased risk of developing such a disease. Furthermore, this invention provides therapeutic and prophylactic methods for treating or preventing Alzheimer's disease and related neurodegenerative disorders using a gene coding for golgin-245. A method of screening for modulating agents of neurodegenerative diseases is also disclosed.

1/14

Figure 1: Identification of Genes Involved in Alzheimer's Disease Pathology



2/14

Figure 2: Identification of differentially expressed genes in a fluorescence differential display screen





Figure 3: Verification of differential expression of golgin-245 by quantitative RT-PCR



4/14

Figure 4: SEQ ID NO. 1; amino acid sequence of golgin-245

Length: 2251 aa

1 MFKKLLKQKIS EEQQQLQQAL APAQASSNSS TPTMRSRSTS SFTEQLDEGT
51 PNRENASTHA SKSPDSVNGS EPSIPQSGDT QSFAQKLQLR VPSVESLFRS
101 PIKESLFRSS SKESLVRTSS RESLNRLDLD SSTASFDFPS DMDSEAEDLV
151 GNSDSLKKEQ LIQRLRRMER SLSSYRGKYS ELVTAYQMLQ REKKKLOGIL
201 SQSQDKSLRR IAELEELQM DQQAQKHLQE EFDASLEEKD QYISVLQTV
251 SLLKQRLRNG PMNVDVLKPL POLEPOAEVF TKEENPESDG EPVVEDGTSV
301 KTLETLOQRV KRQENLLKRC KETIQSHKEQ CTLLTSEKEA LQEQLDERLQ
351 ELEKIKDLHM AEKTKLITQL RDAKNLIEQL EODKGMVIAE TKRQMHEGLE
401 MKEEEEIAQLR SRIKQMTTQG EELREQKEKS ERAAFEELEK ALSTAQKTEE
451 ARRKLKAEMD EQIKTIEKTS EBERISLQQE LSRVKQEVVD VMKKSSEEQI
501 AKLQKLHEKE LARKEQELTK KLQTREREFQ EQMKVALEKS QSEYLKISQE
551 KEQQESLAE ELELQKKAIL TESENKLRLD QQEAETRYRTR ILELESSLEK
601 ELQENKNQSK DLAVHLEAEK NKNKEITVM VEKHKTELES LKHQODALWT
651 EKLQVLKQY QTEMEKLEK CEQKETLLK DKEIIFQAH EEMNEKTEK
701 LDVKQTELES LSSELSEVLK ARHLEELS VLKDQTDKMK QELEAKMDEQ
751 KNHHQQQVDS IIEHEVSIQ RTEKALKDQI NQLELLKER DKHLKEHQAH
801 VENLEADIKR SEGELQQASA KLDVFQSYQS ATHEQTKAYE EQLAQLOQKL
851 LDLETERILL TRQVAEVEAQ KRQVCTELDA HKIQVQDLMO QLEKQNSEME
901 QKVKSILTQVY ESKLEDGNKE QEQTQILVE KENMILQMR ECKKEIEILT
951 QKLSAKEDSI HILNEEYETR FKNQEKMEK VKQKAKEMQE TLKKKLLDQE
1001 AKLKKLENT ALELSQKEKQ FNAKMLEMAQ ANSAGISDAV SRLETNQKEQ
1051 IESLTEVHRR ELNDVISIWE KKLNQQABEL QEIEHQLQE KEQEVAREKQ
1101 KILLFGCEKE EMNKEITWLK EGVKQDITL NELQEQKQK SAHVNSLAQD
1151 ETKLKAHLEK LEVDLNKSLK ENTFLQEQLV ELKMLAEEDK RKVSELTSLK

Proline-rich
domainProline-rich
domain

Granin signature

5/14

1201 KTTDEEFQSL KSSHEKSNKS LEDKSLEFKK LSEELAIQLD ICCKKTEALL
1251 EAKTNELINI SSSKYNAILS RISHCQHRTT KVKEALLIRT CTVSELEAQL
1301 RQLTEEQNTL NISFQQATHQ LEEKENQIKS MKADIESLVT EKEALQKEGG
1351 NQQQAASEKE SCITQLKKEL SENINAVTLM KEELKEKKVE ISSLSKQLTD
1401 LNVQLQNSIS LSEKEAAISS LRKQYDEEKC ELLDQVQDLS FKVDTLSEK
1451 ISALEQVDDW SNKPFSEWKKK AQSRTQHQH TVKELQIQLE LKSKEAYEKD
1501 EQINLLKEEL DQONKRFDCI KGEMEDDKSK MEKKESNLET ELKSQTARIM
1551 ELEDHITQKT IEIESLNEVL KNYNQKDIE HKELVQKLQH FOELGERKDN
1601 RVKEAEKIL TLENQVYSMK AELETKKKEL EHVNLVSKS EELKALED
1651 LESESAAKLA ELKRKAEQKI AAIKKQLLSQ MEEKEEQYK GTESHLSELN
1701 TKLQEREREV HILEEKLKSV ESSQSETLIV PRSAKNVAAY TEQEEADSQG
1751 CVQKTYEEKI SVLQRNLTKE EKLQVRGQE KEETVSSHFE MRCQYQERLI
1801 KLEHAEAKQH EDQSMIGHLQ EELEEKNNKY SLIVAQHVEK EGGKNNIQAK
1851 QNLENVFDDV QKTLQEKELT CQILEQKIKE LDSCLVRQKE VHRVEMEELT
1901 SKYEKLQALQ QMDGRNKPTL LLEENTEEKS KSHLVQPKLL ENMEAQHNDL
1951 EFKLAGAERE KQKLGKEIVR LQKDLRMLRK EHQQELEILK KEYDQEREEK
2001 IKQEQEDLEL KHNSTLKQIM REFNTQLAQK EQELEMTIKE TINKAQVEA
2051 ELLESHQEET NOLLKKIAEK DDDLKRTAKR YEEILDAREE EMTAKVRDLQ
2101 TQLEELQKKY QOKLEQENB GNDMVTIMEL QTQLAQKTTL ISDSKLKEQE
2151 FREQIHNL ED RLKKYKENVY ATTVGTPYKG GNLYHTDVSL FGEPTEFEYL
2201 RKVLFEYMMG RETKIMAKVI TTVLKFPDDQ TOKILEREDA RLMSWLRSSE
2251 *

GRIP-domain and
highly conserved
tyrosine residue

Figure 5: SEQ ID NO: 2

Length: 7743 January 9, 2002 17:30 Type: N Check: 12 ..

1 GCAACGAAGG TACCATGGCC GTTGTCTGCG CCGCCGCGGC TCCCAGGGCT
51 GGATGGGGGG CCGAGGCCAG CCAGTGGCAC CCGGAAGAAA GAGACGCGGC
101 GCGGCGGACG CCGACACCTT CAGGACGAGT GTCCGGACTT GCCCAGAGCC
151 TCAAGGAGGA GACGGCGAGG CCGGGCCCCC GCTGTCCCTG GTGTAAAGAA
201 GTCGCCGTAG CCGTCGCGGC CGGGACTCCC CGGGCTCTCG CCCTTCAGGT
251 TTCGTTGACA CTCAGGACCG TACGTACGCT GCGCCATGTT CAAGAAACTG
301 AAGCAAAGA TCAGCGAGGA GCAGCAGCAG CTCCAGCAGG CGCTGGCTCC
351 TGCTCAGGCG TCCTCCAATT CTTCAACACC AACAGAATG AGGAGCAGGA
401 CATCTTCATT TACAGAGCAA CTTGATGAAG GTACACCCAA TAGAGAGAAT
451 GCATCTACTC ATGCCTCGAA ATCTCTGAC AGTGTTAATG GAAGTGAACC
501 AAGCATTCTT CAGTCAGGTG ACACACAGTC TTTTGCACAG AAGCTCCAGC
551 TCCGGGTGCC CTCCGTGGAG TCTTTGTTTC GAAGTCCGAT AAAGGAATCT
601 CTATTCCGGT CTTCTTCTAA AGAGTCTTTG GTACGAACAT CTTCCAGAGA
651 ATCCCTGAAT CGACTTGACC TGGACAGTTC TACTGCCAGT TTTGATCCAC
701 CCTCTGATAT GGATAGCGAG GCTGAAGACT TGGTAGGGAA TTCAGACAGT
751 CTCAACAAAG AACAGTTGAT TCAGCGGTTG CGAAGAATGG AACGAAGCTT
801 AAGTAGCTAC AGGGGAAAAT ATTCTGAGCT TGTACAGCT TATCAGATGC
851 TTCAGAGAGA GAAGAAAAG CTACAAGGTA TATTAAGTCA GAGTCAGGAT
901 AATCACTTC GGAGAATAGC AGAATTAAGA GAGGAGCTCC AATGGACCA
951 GCAGGCAAAG AAACATCTGC AAGAGGAGTT TGATGCATCT TTAGAGGAGA
1001 AAGATCAGTA TATCAGTGT CTCCAAACTC AGGTTTCTCT ACTGAAACAA
1051 CGATTACGAA ATGGCCCGAT GAATGTTGAT GTACTGAAAC CACTTCCTCA
1101 GCTGGAACCA CAGGCTGAAG TCTTCACTAA AGAAGAGAAT CCAGAAAGTG
1151 ATGGAGAGCC AGTAGTGAA GATGGAACCT CTGTAAAAAC ACTGGAACA
1201 CTCCAGCAA GAGTGAAGCG TCAAGAGAAC CTACTTAAGC GTTGTAAAGGA
1251 AACAAATCAG TCACATAAGG AACAAATGTAC ACTATTAACT AGTGAAAAAG

7/14

1301 AAGCTCTGCA AGAACAACTG GATGAAAGAC TTCAAGAACT AGAAAAGATA
1351 AAGGACCTTC ATATGGCCGA GAAGACTAAA CTTATCACTC AGTTGCGTGA
1401 TGCAAAGAAC TTAATTGAAC AGCTTGAACA AGATAAGGGA ATGGTAATCG
1451 CAGAGACAAA ACGTCAGATG CATGAAACCC TGGAAATGAA AGAAGAAGAA
1501 ATTGCTCAAC TCCGTAGTCG CATCAAACAG ATGACTACCC AGGGAGAGGA
1551 ATTACGGGAA CAGAAAGAAA AGTCCGAAAG AGCTGCTTTT GAGGAACTTG
1601 AAAAAGCTTT GAGTACAGCC CAAAAACAG AGGAAGCACG GAGAAAAC TG
1651 AAGGCAGAAA TGGATGAACA AATAAAACT ATCGAAAAA CAAGTGAGGA
1701 GGAACGCATC AGTCTTCAAC AGGAATTAAG TCGGGTGAAA CAGGAGGTTG
1751 TTGATGTAAT GAAAAATCC TCAGAAGAAC AAATTGCTAA GCTACAGAAG
1801 CTTCATGAAA AGGAGCTGGC CAGAAAAGAG CAGGAACTGA CCAAGAAGCT
1851 TCAGACCCGA GAAAGGGAAT TTCAGGAACA AATGAAAGTA GCTCTTGAAA
1901 AGAGTCAATC AGAATATTG AAGATCAGCC AAGAAAAAGA ACAGCAAGAA
1951 TCTTTGGCCC TAGAAGAGTT AGAGTTGCAG AAAAAAGCAA TCCTCAGAGA
2001 AAGTGAAAAT AACTTTCGGG ACCTTCAGCA AGAAGCAGAG ACTTACAGAA
2051 CTAGAATTCT TGAATTGGAA AGTTCTTTGG AAAAAAGCTT ACAAGAAAAC
2101 AAAATCAGT CAAAAGATT GGCTGTTTAT CTGGAAGCTG AAAAAATAA
2151 GCACAATAAG GAGATTACAG TCATGGTTGA AAAACACAAG ACAGAATTGG
2201 AAAGCCTTAA GCATCAGCAG GATGCCCTTT GGACTGAAAA ACTCCAAGTC
2251 TTAAAGCAAC AATATCAGAC TGAAATGGAA AACTTAGGG AAAAGTGTGA
2301 ACAAGAAAAA GAAACATTGT TGAAAGACAA AGAGATTATC TTCCAGGCCC
2351 ACATAGAAGA AATGAATGAA AAGACTTTAG AAAAGCTTGA TGTGAAGCAA
2401 ACAGAACTAG ATCATTATC TTCTGAACTG TCAGAAGTAT TAAAGCCCCG
2451 TCACAAACTA GAAGAGGAAC TTTCTGTTCT GAAAGATCAA ACAGATAAAA
2501 TGAAGCAGGA ATTAGAGGCC AAGATGGATG AACAGAAAAA TCATCACCAG
2551 CAGCAAGTTG ACAGTATCAT TAAAGAACAC GAGGTATCTA TCCAGGGGAC
2601 TGAGAAGGCA TTAAAGATC AAATTAATCA ACTTGAGCTT CTCTTGAAGG
2651 AAAGGGACAA GCATTTGAAA GAGCATCAGG CTCATGTAGA AAATTTAGAG
2701 GCAGATATTA AAAGGTCTGA AGGGGAACTC CAGCAGGCAT CTGCTAAGCT
2751 GGACGTTTTT CAGTCTTACC AGAGTGCCAC ACATGAGCAG ACAAAGCAT
2801 ATGAGGAACA GTTGGCCCAA TTGCAGCAGA AGTTGTTGGA TTGGAAACA

8/14

2851 GAAAGAATTC TTCTTACCAA ACAGGTTGCT GAAGTTGAAG CACAAAAGAA
2901 AGATGTTTGT ACTGAGTTAG ATGCTCACAA AATCCAGGTG CAGGACTTAA
2951 TGCAGCAACT TGAAAAACAA AATAGTGAAA TGGAGCAAAA AGTAAAATCT
3001 TTAACCCAAG TCTATGAGTC CAACTTGAA GATGGTAACA AAGAACAGGA
3051 ACAGACAAAG CAAATCTTGG TGGAAAAGGA AAATATGATT TTACAAATGA
3101 GAGAAGGACA GAAGAAAGAA ATTGAGATAC TCACACAGAA ATTGTCAGCC
3151 AAGGAGGACA GTATTCATAT TTTGAATGAG GAATATGAAA CCAAATTTAA
3201 AAACCAAGAA AAAAAGATGG AAAAAGTTAA GCAGAAAGCA AAGGAGATGC
3251 AAGAAACGTT AAAGAAAAAA TTACTGGATC AGGAAGCCAA ACTTAAGAAA
3301 GAGCTTGAAA ATACTGCTCT AGAGCTTAGT CAGAAAGAAA AACAGTTTAA
3351 TGCCAAAATG CTGGAAATGG CACAGGCTAA CTCAGCTGGA ATCAGTGATG
3401 CAGTGTCAAG ACTGGAAACA AACC AAAAAG AACAAATAGA AAGTCTTACT
3451 GAGGTTTCATC GACGAGAACT CAATGATGTC ATATCAATCT GGGAAAAGAA
3501 ACTTAATCAG CAAGCTGAAG AACTTCAGGA AATACATGAA ATCCAATTAC
3551 AGGAAAAAGA ACAAGAGGTA GCAGAACTGA AACAAAAGAT CCTCCTATTT
3601 GGGTGTGAAA AAGAAGAGAT GAACAAGGAA ATAACATGGC TGAAGGAAGA
3651 AGGTGTTAAG CAGGATACAA CATTAAATGA ATTACAGGAA CAGTTAAAGC
3701 AGAAGTCTGC CCATGTGAAT TCTCTTGCAC AAGATGAAAC TAACTGAAA
3751 GCTCATCTTG AAAAGCTAGA GGTGACTTG AATAAGTCTC TGAAGGAAAA
3801 TACTTTTCTT CAAGAGCAGC TAGTTGAACT GAAGATGCTG GCAGAAGAAG
3851 ATAAGCGGAA GGTCTCTGAG TTGACTAGCA AGTTGAAAAC CACAGATGAA
3901 GAATTCAGAA GTTTGAAATC TTCACATGAA AAAAGTAACA AAAGCCTAGA
3951 GGACAAGAGC TTGGAATTTA AAAAAGTCTC TGAGGAACTA GCGATTCAGC
4001 TAGATATTTG CTGTAAGAAA ACCGAAGCCT TATTAGAAGC TAAAACAAAT
4051 GAGCTAATCA ACATTAGTAG TAGTAAACT AATGCCATTC TTTCTAGGAT
4101 TTCTCATTGT CAGCACCCTA CAACTAAAGT TAAGGAGGCA CTGTTAATTA
4151 AAAGTGCAC AGTTTCTGAA TTAGAAGCAC AACTTAGACA GTTGACAGAG
4201 GAGCAAAATA CACTAAATAT TTCTTTTCAA CAGGCTACTC ATCAGTTAGA
4251 AGAAAAAGAA AATCAAATTA AGAGCATGAA GGCTGATATT GAAAGTCTTG
4301 TAACAGAAAA AGAAGCCTTA CAGAAGGAAG GAGGCAATCA GCAACAGGCT
4351 GCTTCTGAAA AGGAGTCTTG TATAACACAG TTGAAGAAAG AGTTATCTGA
4401 AAACATCAAT GCTGTCACAT TGATGAAAGA AGAGCTTAAA GAAAAAAG

4451 TTGAGATTAG CAGTCTTAGT AAACAACATA CTGATTTGAA TGTTTCAGCTT
 4501 CAAATATAGCA TCAGCCTATC CGAAAAAGAA GCAGCCATTT CATCACTAAG
 4551 AAAGCAGTAT GATGAAGAAA AATGTGAATT GCTGGATCAG GTGCAAGATT
 4601 TATCTTTTAA AGTTGACACT CTGAGTAAAG AGAAAATTC TGCTCTTGAG
 4651 CAGGTAGATG ACTGGTCCAA TAAATTCTCA GAATGGAAGA AGAAGCACA
 4701 GTCAAGATTT ACACAGCATC AAAACACTGT TAAAGAATTG CAGATCCAGC
 4751 TTGAGTTAAA ATCAAAGGAA GCTTATGAAA AGGATGAGCA GATAAATTTA
 4801 TTGAAGGAAG AGCTTGATCA GCAAAATAAA AGATTTGATT GTTTAAAGGG
 4851 TGAAATGGAA GACGACAAGA GCAAGATGGA GAAAAAGGAG TCTAATTTAG
 4901 AAACAGAGTT AAAGTCTCAA ACAGCAAGAA TTATGGAATT AGAGGACCAT
 4951 ATTACCCAGA AACTATTGA AATAGAGTCC TTAAATGAAG TTCTTAAAAA
 5001 TTACAATCAA CAAAAGGATA TTGAACACAA AGAATTGGTT CAGAACTTC
 5051 AACATTTTCA AGAGTTAGGA GAAGAAAAGG ACAACAGGGT TAAAGAAGCT
 5101 GAAGAAAAAA TCTTAACACT TGAAAACCAA GTTTATTCCA TGAAAGCTGA
 5151 ACTTGAAACT AAGAAGAAAG AATTAGAACA TGTGAATTTA AGTGTGAAAA
 5201 GCAAAGAGGA GGAGTTAAAG GCATTGGAAG ATAGGCTTGA GTCAGAAAGT
 5251 GCTGCAAAAT TAGCAGAGTT GAAGAGAAAA GCTGAACAAA AAATTGCTGC
 5301 CATTAAGAAG CAGTTGTTAT CTCAAATGGA AGAGAAAGAA GACAGTATA
 5351 AAAAAGGTAC AGAAAGCCAT TTGAGTGAGC TAAATACAAA ATTGCAGGA
 5401 AGAGAAAGGG AAGTTCACAT CTTGGAAGAA AACTTAAAGT CAGTGGAAAG
 5451 TTCACAGTCA GAAACATTAA TTGTACCCAG ATCAGCAAAA AATGTGGCAG
 5501 CATATACTGA ACAAGAAGAA GCAGATTCCC AAGGCTGTGT GCAGAAGACA
 5551 TATGAAGAAA AAATCAGTGT TTTACAAAGA AACTTAACTG AAAAAGAAAA
 5601 GCTATTGCAG AGGGTAGGGC AGGAAAAAGA AGAGACAGTT TCTTCTCAT
 5651 TTGAAATGCG ATGCCAATAC CAGGAGCGCT TAATAAAGCT AGAACATGCT
 5701 GAGGCAAAGC AACATGAAGA TCAAAGTATG ATAGGTCATC TTCAAGAGGA
 5751 GCTTGAAGAA AAAACAAGA AATATTCCTT GATAGTAGCC CAGCATGTGG
 5801 AAAAAGAAGG AGGTAAAAAT AACATACAGG CAAAGCAAAA CTTGGAAAT
 5851 GTGTTTGACG ACGTCCAGAA AACCTCCAG GAGAAGGAAC TAACCTGTCA
 5901 GATTTTGGAG CAAAAGATAA AAGAGCTGGA TTCCTGCTTA GTAAGACAGA
 5951 AAGAAGTACA TAGAGTTGAA ATGGAAGAGT TGACCTCAA ATATGAAAAA

10/14

6001 TTACAGGCTT TACAACAGAT GGATGGAAGA AATAAACCCA CAGAACTTTT
6051 GGAAGAAAAG ACTGAAGAAA AGTCCAAATC ACATTGGTC CAACCCAAAT
6101 TGCTTAGTAA CATGGAAGCC CAGCACAATG ATCTGGAGTT TAAATTAGCC
6151 GGGGCAGAAC GGGAGAAACA GAACTGGGC AAGGAGATTG TTAGATTGCA
6201 GAAAGACCTT CGAATGTTGA GAAAGGAGCA TCAGCAAGAA TTGGAAATAC
6251 TAAAGAAAGA ATATGATCAA GAAAGGGAAG AGAAATCAA ACAGGAGCAG
6301 GAAGATCTTG AACTGAAGCA CAATTCCACA TTAAACAGC TGATGAGGGA
6351 GTTTAATACA CAGCTGGCAC AAAAGGAACA AGAGCTGGAA ATGACCATAA
6401 AAGAACTAT CAATAAGGCC CAGGAGGTGG AGGCTGAACT TTTAGAAAGC
6451 CATCAAGAAG AGACAAATCA GTTACTTAAA AAAATTGCTG AGAAAGATGA
6501 TGATCTAAAA CGAACAGCCA AAAGATATGA AGAAATCCTT GATGCTCGTG
6551 AAGAAGAAAT GACTGCAAAA GTAAGGGACC TGCAGACTCA ACTTGAGGAG
6601 CTGCAGAAGA AATACCAGCA AAAGCTAGAG CAGGAGGAGA ACCCTGGCAA
6651 TGATAATGTA ACAATTATGG AGCTACAGAC ACAGCTAGCA CAGAAGACGA
6701 CTTTAAATCAG TGATTGAAA TTGAAAGAGC AAGAGTTCAG AGAACAGATT
6751 CACAATTTAG AAGACCGTTT GAAGAAATAT GAAAGAATG TATATGCAAC
6801 AACTGTGGGG ACACCTTACA AAGGTGGCAA TTTGTACCAT ACGGATGTCT
6851 CACTCTTTGG AGAACCTACC GAATTGAGT ATTTGCGAAA AGTGCTTTTT
6901 GAGTATATGA TGGGTCGTGA GACTAAGACC ATGGCAAAG TTATAACCAC
6951 CGTACTGAAG TTCCCTGATG ATCAGACTCA GAAATTTTG GAAAGAGAAG
7001 ATGCTCGGCT GATGTTTACT TCACCTCGCA GTGGTATCTT CTGAGTAAAC
7051 CATCAGTCTG TGCTTAGTTA ACATGTGTCA TGGCTCCGAT CTTCATCTTG
7101 AAGAAGAGTG ACATTGGGTG ACTGCTGCTT GGAAACTGT CCACACTTGC
7151 TACTCTTTGA GAATGAAGTT GTCATTGAGG GCCCCTCATG TAGCCAAAAG
7201 ACCAAGAAAA ATCTGGCCCA CAGATAAGTT GCAGACTGCC TTTAAATAG
7251 ATTTTATCAG TGGAGAAATG GTGATAGTTT TTTCTTCAGT TTTCTCTTGG
7301 GAAGAGTTTT ATGTTGTTA AAAGATATTT TGATAACTTA ACCTGCTTTA
7351 TGGGCTTACA TAATATTCCT TTCATCCATT CTTTTTAAAG AACGGCTTAC
7401 CTTTCCTATT TATTTTLAGG GTGATTTTTT AAAAGACTT GTGCAATACA
7451 TTTTGAGGTG AACTTAGTG GATTTTTTCT GATAAATTAG AGCATTTAAT
7501 TGAATTTTT ATTGAGTTG ATCTGTTGAA TATTTGCTAA AGACCAAGTC
7551 TTTAAGCTAA GACATGTAAA AATCCCAA TGGCAGTACC TCATTGTTTA



11/14

7601 CTTAGCTTTT GTACTTATAT TTTTCAGAGG AAAAAACACT ACTGTAAATT

7651 GTGAATAGCC AATACATAAC TGTATTGTAT GCAAACTCTGT GATTGTTGGC

7701 AGTGTCATCT CTGAGAAACA GATAAATAAA GTTTATTTAC TAT

12/14

Figure 6: SEQ ID NO. 3

Length: 36 bp

1 AGTTAAGTTT CTTTGTA AAA CACTGATTTT TTCTCC

13/14

**Figure 7: Alignment of SEQ ID NO. 3
with SEQ ID NO. 2, human golgin-
245 cDNA (GenBank accession
number U41740)**

```
36 GGAGAAAAAATCAGTGTTTACAAAGAACTTAACT 1
   | | | | | | | | | | | | | | | | | | |
5554 GAAGAAAAAATCAGTGTTTACAAAGAACTTAACT 5589
```

14/14

Table 1

sample	Δ (fold) (temporal/frontal cortex)
patient 1	0.98
patient 2	1.55
patient 3	1.56
patient 4	1.30
patient 5	1.32
patient 6	2.91
patient 7	1.06
control 1	0.86
control 2	0.86
control 3	1.00
control 4	0.87

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☒ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☒ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☒ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.